

# Bipolar disorder in pregnancy

## A guide for women and their families



Bipolar disorder is a mental health condition that causes mood shifts, usually between episodes of depression and mania (extremely high mood). The risk of developing bipolar disorder or having a relapse increases during pregnancy.

### How common is bipolar disorder?

Bipolar disorder is experienced by 1 in 100 people in the Australian population.

### What causes bipolar disorder?

While the cause of bipolar disorder is unknown, it is often inherited or linked to stressful life events.

### What are the signs and symptoms?

#### Common behaviour associated with depression

- Moodiness that is out of character
- Increased irritability and frustration
- Finding it hard to take minor personal criticisms
- Spending less time with friends and family
- Loss of interest in food, sex, exercise or other pleasurable activities
- Being awake throughout the night
- Increased alcohol and drug use
- Staying home from work or school
- Increased physical health complaints like fatigue or pain
- Slowing down of thoughts and actions.

#### Common behaviour associated with mania

- Increased energy
- Irritability
- Overactivity
- Increased spending
- Being reckless or taking unnecessary risks
- Increased sex drive
- Racing thoughts
- Rapid speech
- Decreased sleep
- Grandiose ideas
- Hallucinations and/or delusions.

### Support in the antenatal period

Attending antenatal care is important if you have bipolar disorder because relapse is common during pregnancy, especially if you have stopped taking medications. Your doctor or midwife can also give you information about nutrition and ceasing smoking, illicit substance use and alcohol intake in pregnancy.

### How is bipolar disorder treated?

#### Talking therapies

Talking therapies can be very helpful in treating symptoms of depression and anxiety, which are common in people with bipolar disorder. They can help you recognise and change the negative thinking and feelings that depression brings. They also give you useful tools to stop anxiety coming back.

Talking therapies include:

- cognitive-behavioural therapy (CBT): the cognitive (or thinking) part of this therapy teaches you to think logically and challenge negative thoughts. The behaviour part helps you change the way you react in situations and can help you to get involved in activities that you have been avoiding or have stopped doing
- interpersonal therapy (IPT): helps you to find new ways to connect with others, and overcome losses, challenges and conflicts that you may have.

Registered practitioners with appropriate training and experience provide talking therapies for individuals or groups.

#### Medicines

The main medicines used to treat bipolar disorder are called mood stabilisers. These help to reduce the likelihood of your symptoms recurring (relapse). The most common mood stabiliser is lithium, which can be safely used during pregnancy. Other mood stabilisers that may be used are carbamazepine and lamotrigine. Sodium valproate is not safe for use in pregnancy.



# Bipolar disorder in pregnancy

## A guide for women and their families

Depending on your symptoms, your doctor may also prescribe other medicines.

- Antidepressants are used to treat symptoms of depression and/or anxiety. Antidepressants that can be safely used during pregnancy are called selective serotonin reuptake inhibitors (SSRIs). Your doctor may also suggest another type of antidepressant called tricyclic antidepressants, especially if they have worked for you in the past.
- Antipsychotics help improve problems of mood, thinking and socialising and anxiety or agitation. Other than clozapine, antipsychotics can be safely used during pregnancy. As some antipsychotics increase weight gain, it is good to talk to your doctor or another health professional about managing your weight if you are taking antipsychotics.

It is important not to change or stop taking your medicines without talking to your doctor.

## Advice for women experiencing bipolar disorder in pregnancy

### Seek help and treatment

- Seek help from a doctor or other health professional.
- Learn about effective treatments.
- Call a support service or mental health crisis line if other help is not available.

### Accept help and support

- Develop a support system of friends, family and professionals and accept help.
- Discuss your feelings with your partner, family and friends.

### Look after your physical health

- Try to eat healthy meals, including fruit, vegetables, whole grains and lots of water.
- Plan some enjoyable physical activity every day.
- Try to establish good sleeping patterns.
- Practice techniques to reduce stress, such as muscle relaxation and deep breathing.

## Advice for family and friends providing support

### Listen and reassure:

- Encourage the woman to discuss her symptoms.
- Explain that bipolar disorder can be treated and managed.

### Provide information:

- Give the woman good quality information about bipolar disorder in pregnancy, such as this fact sheet.
- Give details of helplines if she is feeling distressed and needs support.
- Offer information to the woman's partner/others.

### Direct to care and support:

- Encourage the woman to talk to her GP or other health professional.
- Encourage the woman to identify and draw on possible supports and services that may be available to her for practical and/or emotional support.
- Remind the woman that she can go to her doctor or local hospital if she is at risk of harming herself or others.

